

POST OPERATIVE INFORMATION

Carpal Tunnel Decompression

Normal Symptoms:

- · Discomfort in wrist
- Mild swelling

The following symptoms are **NOT** normal:

- Increased swelling
- Redness around the incision
- Excessive bleeding
- · Fever of chills

If you experience any of the above symptoms you should seek medical advice. Please contact your nearest Emergency Department.

Diet:

Normal healthy diet as tolerated.

Activity:

Avoid heavy lifting.

Medications:

- Take medications as directed by your Doctor.
- Regular Panadol may be useful for pain.
- Cold pack applied at regular intervals can help reduce post-operative swelling.

Driving:

Be guided by your pain and avoid driving for one week.

Dressings / Wound care:

- Do not get the dressing wet.
- Keep dressings on as directed by Doctor.

Hygiene:


You may shower as desired.

Legal requirements following an anaesthetic or sedation:

For a 24 hour period you are <u>NOT</u> allowed to:

- · Drive a car as directed by Doctor
- · Operate machinery
- · Drink alcohol
- Sign any legal or important documents.

Follow-up Appointment





DR L. ANDREW ASHTON

B.A., B.Med., FRACS (Orth) Orthopaedic Surgeon

E secretary@oso.com.au T 0263 601 500 309 Anson Street, Orange NSW 2800 PO Box 8136, East Orange 2800 www.orthopaedicsurgeonorange.com.au

PROVIDER NO. 204803DF **ABN** 13371125435